

Effects of disinfection on the molecular detection of porcine epidemic diarrhea virus

Bowman, A.; Nolting, J.; Nelson, S.; Bliss, N.; Stull, J.; Wang, Q.; Premanandan, C.

ABSTRACT

Routine detection of porcine epidemic diarrhea (PEDv) is currently limited to RT-PCR testing, as it is the only test method that can directly detect PEDv. Because RT-PCR only detects the viral RNA, a positive RT-PCR result only indicates the presence of PEDv viral RNA, but does not mean viable and infectious virus is present. Accelerated Hydrogen Peroxide®(AHP®) is a relatively new yet proven technology that is capable of disinfecting PEDv but may still leave inactivated RNA strands on surfaces, and therefore has a history of producing RT-PCR positive test results. In this study AHP was tested along with a number of other disinfectant actives as agents against PEDv using RT-PCR. Positive RT-PCR results were tested to show how AHP was able to fully inactivate any remaining RNA on the surface. Therefore AHP can be used as an alternative disinfectant that is effective against PEDv without the negative toxicity, environmental, safety and compatibility profiles.

BACKGROUND

Contaminated transportation equipment has been linked to the spread of several other important swine diseases making trailer disinfection common among pork producers. Efficient disinfection for PEDv in animal contact spaces, including trailers is one of the primary methods used to control the spread of disease. Due to the limited testing options and the implications of environmental contamination, individuals are using RT-PCR to test trailers following disinfection to ensure that the equipment is free of PEDv before contact with animals. RT-PCR tends to underestimate disinfection efficacy compared to infectivity assays; meaning, RT-PCR positive results are obtained when in fact the trailer has been effectively disinfected.

STUDY

The purpose of this study was to examine the effect of disinfectants on RT-PCR results for PEDv and explore the practical solutions to produce RT-PCR negative trailers after they have been contaminated with PEDv. Five classes of disinfectants, including AHP, were evaluated at varying concentrations, both in the presence and absence of swine feces. No infectious PEDv was recovered after treatment with evaluated disinfectants. Additionally, all tested disinfectants,

except for 0.17% sodium hypochlorite dramatically reduced RT-PCR values. However, no disinfectants eliminated RT-PCR detection of PEDv across all replicates; although 0.52%, 1.03%, and 2.06% solutions of sodium hypochlorite and 0.5% AHP did produce intermittently RT-PCR negatives. To simulate field conditions in a second attempt, PEDv was applied to pitted aluminum trays, which were treated with either 2.06% sodium hypochlorite or 0.5% AHP. Post-treatment surface swabs of the trays tested RT-PCR positive but were not infectious to cultured cells or naive pigs. Ultimately, viable PEDv was not detected following application of each of the tested disinfectants, even though in most cases RT-PCR detection of viral RNA remained.

CONCLUSION

Because PEDv strains are difficult, time consuming and expensive to test using cell culture methods, the pork industry is relying upon RT-PCR for testing. Ultimately, all the tested disinfectants, including AHP were able to inactivate PEDv but few prevented RT-PCR detection of the Viral RNA. This study also indicates that the use of RT-PCR as a method to indicate the presence of infective PEDv on a surface is not reliable. One must take caution if using sodium hypochlorite as it is known to cause topical chemical burns, respiratory irritation and pulmonary edema, and can cause corrosion of metals and deterioration of rubber objects. AHP can therefore be used as an alternative disinfectant that is effective against PEDv, without the negative toxicity, environmental, safety and compatibility profiles.

IMPLICATIONS FOR AHP

AHP Disinfectants carry claims against Porcine Epidemic Diarrhea (PEDv)

- All AHP products have been tested and are effective against both enveloped viruses (easy to kill) such as PEDv and non-enveloped viruses (hard to kill) such as Parvovirus

AHP Disinfectants have realistic contact times

- Short contact times ensure surfaces remain wet for the required contact time, providing comfort and confidence that disinfection has occurred
- AHP has been tested when mixed with propylene glycol to ensure the product does not freeze and has been proven effective in freezing temperatures



AHP Disinfectants provide the perfect balance between safety and efficacy

- All AHP products at in-use solutions do not require the use of Personal Protective Equipment (PPE) without sacrificing germicidal potency against pathogens such as PEDv

AHP Disinfectants are One-Step Disinfectant Cleaners

- One-Step Disinfectant Cleaners provide you with added confidence - you clean while you disinfect, and disinfect while you clean

AHP Disinfectants are environmentally sustainable

- AHP's active ingredient, hydrogen peroxide, breaks down into water and oxygen leaving no active residues
- AHP products have been inherently proven to be biodegradable indicating that properly maintained septic systems and/or manure pits can handle normal wastes generated from the use of AHP formulations

REFERENCE

Bowman, A. et al. (2015). *Effects of disinfection on the molecular detection of porcine epidemic diarrhea virus*. *Vet. Microbiol.* <http://dx.doi.org/10.1016/j.vetmic.2015.05.027>